

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number
WO 02/22590 A1

(51) International Patent Classification⁷: C07D 277/82 //
C07M 7:00

(74) Agent: DUXBURY, Stephen; Arnold & Siedsma, Sweel-
inckplein 1, NL-2517 GK The Hague (NL).

(21) International Application Number: PCT/NL01/00680

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date:
14 September 2001 (14.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
200002262 18 September 2000 (18.09.2000) ES

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): SYN-
THON B.V. [NL/NL]; Microweg 22, NL-6545 CM
Nijmegen (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): POSPISILIK,
Karel [CZ/CZ]; Jezkova 1, CZ-63800 Brno (CZ).
HOORN, Hans, Jan [NL/NL]; Hegdambroek 17-24,
NL-6546 VX Nijmegen (NL). PETERS, Theodorus,
Hendricus, Antonius [NL/NL]; G.A. van Nispenstraat 22,
NL-6814 JB Arnhem (NL). LEMMENS, Jacobus, Maria
[NL/NL]; Lodewijkstraat 30, NL-6585 CM Mook (NL).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR PREPARATION OF 2-AMINO-6-(ALKYL) AMINO-4,5,6,7-TETRAHYDROBENZOTHAZOLES

(57) Abstract: A new process to obtain pramipexole and related products is described. The process involves the reaction of new compounds of formula (6), wherein R is hydrogen or acyl group, R₃ and R₄ are either the same and each of them represents an alkoxy group of 1-4 carbons or they together form a C₂-C₅ alkylenedioxy group or an oxo-group, with an alkylamine in the presence of a reducing agent or a hydrogen gas with hydrogenation catalyst. A process to obtain new compounds of formula (6) is also described.




WO 02/22590 A1

PROCESS FOR PREPARATION OF 2-AMINO-6-(ALKYL)AMINO-4,5,6,7-TETRAHYDROBENZOTHAZOLES

5 The present invention relates to a novel process for preparing 2-amino-6-alkylamino-4,5,6,7-tetrahydrobenzothiazoles and to certain intermediates useful therein.


Background and prior art

10 The 2-(acyl)amino-6-(substituted)amino-4,5,6,7-tetrahydrobenzothiazoles having the general formula (1):



15 wherein R is hydrogen or acyl group, R₁ is hydrogen, alkyl or aralkyl group and R₂ is hydrogen,


are useful pharmaceutical agents. Some of these compounds are known to have dopamine D-2 agonist activity. The compounds of formula (1) include S(-)-2-amino-6-propylamino-4,5,6,7-tetrahydrobenzothiazole (pramipexole) of the formula (2)



which is a commercial product used for treatment of Parkinson's disease and schizophrenia and is marketed, in a form of a dihydrochloride, under several brand names e.g. Mirapexin[™].

The compounds of formula (1) are described in EP 186087.

5 Several methods for preparing compounds of the above formula (1) are suggested in EP 186087 and EP 207696. A common core of these methods is a process comprising ring halogenation (preferably bromination) of a substituted aminoketone (3) and the condensation of the so obtained alpha-halogenaminoketone (4) with thiourea or N-acylthiourea to form a 2-aminotetrahydrobenzothiazole ring, as shown in the following
10 scheme:



15 Dependent on the nature of substituents R₁, R₂ in (3) and (4) and on the desired structure of the product (1), the corresponding amino substituents which are desired to be in position 6 of compounds (1) may or have to be accordingly modified before and after this two-step condensation.

20 A compound of formula (1), wherein both R₁ and R₂ are hydrogen, is prepared from a compound (3) wherein either R₁ is an amino-protective group such as an acyl or alkoxycarbonyl group and R₂ is hydrogen or R₁-R₂ together form an imino-protective group such as phthalimidogroup. After halogenation and condensation with thiourea, the protective group is removed in a separate step.

A compound of formula (1), wherein R₁ is acyl and R₂ is hydrogen, is prepared from a compound (3) wherein R₁ is acyl and R₂ is hydrogen.

25 A compound of formula (1), wherein R₁ is alkyl or aralkyl and R₂ is hydrogen, is either prepared from a compound (3) wherein R₁ is alkyl or aralkyl and R₂

is hydrogen or a protective group (with subsequent deprotection in that later case), or it can be prepared by alkylation /aralkylation of the compound (1) wherein both R_1 and R_2 is hydrogen or, finally, it can be prepared by metal hydride or borane reduction of the acyl group in a compound of formula (1) wherein R_1 is acyl or arylacyl and R_2 is hydrogen .

In all the above cases, the acyl substituent R should be furthermore hydrolysed to hydrogen, whenever necessary.

If desired, the produced compounds of formula (1) may be converted into salts with inorganic and organic acids, particularly with acids which are pharmaceutically acceptable.

Thus, in total, the overall synthetic process represents a sequence of at least four synthetic steps. In practice, a primary aminogroup or a secondary alkylaminogroup present in the compounds of formula (1) cannot be introduced and maintained during oxidation, bromination and cyclization step without introduction a protective group, due to its reactivity. The protective group must be removed afterwards.

In the case of pramipexole and similar compounds, the following synthetic sequence from the cited prior art is suggested to be most useful .

The last step of the overall synthesis comprises reductive alkylation of 2,6-diamino-4,5,6,7-tetrahydrobenzothiazole (compound (1), $R_1=R_2=H$), e.g. by propionaldehyde/sodium borohydride; the starting diamino-compound has been prepared by a sequence starting from 4-aminocyclohexanol which has been acetylated or phthalidated and subsequently oxidised to yield acetamido- or 4-phthalimidocyclohexanone which has been monobrominated and subsequently reacted with thiourea to give 6-acetamido (or 6-phthalimido)-2-amino-4,5,6,7-tetrahydrobenzothiazole (compound (1), $R_1=H$, $R_2=CO-CH_3$ and/or $R_1, R_2=phthalimido$); finally, the protective acetyl or phthalimido group has been removed.

Thus, in practice, the overall synthetic sequence, starting from commonly available material, represents six synthetic steps.


In addition, the compounds of formula (1) have an asymmetric carbon and they exist either as single enantiomers and/or in a racemic form. The pharmacological activity of compounds of formula (1) is however generally connected only or mainly with one stereoisomer thereof; for instance, pramipexole is marketed as a single S(-) isomer and the dopaminergic activity of the said isomer is twice as high as that of the R(+) isomer. The cited prior art process allows preparing only a racemate. It is anticipated that, if the product of formula (1) has a chiral atom, the produced racemic compound may be resolved into optical isomers by classical methods such as chromatography or fractional crystallisation. Generally it should thus be expected that, in industrial scale, the produced racemic compounds of formula (1) are resolved into optical isomers by adding a next production sequence comprising steps of forming a salt with an appropriate optically active acid, resolution of salts by fractional crystallisation and, if necessary, liberating the free base of the resolved product from the salt. An example of such resolution process in the case of producing optically pure pramipexole has been disclosed by Schneider and Mierau in J.Med.Chem 30, 494 (1987), using the diaminoderivative (compound (1), $R=R_1=R_2=H$) as a substrate and L(+) tartaric acid as a resolution agent. Following to the resolution, optically active pramipexole has been prepared by two-step propylation of the single enantiomer of the diamino-precursor comprising reaction with propionanhydride followed by reduction of the propionyl intermediate.

It is apparent that the prior art processes for preparing compounds of formula (1) suffers from severe drawbacks, as they are lengthy and economically undesirable. Thus, there exist a need for a more straightforward production process.

Description of the invention

As the result of the present inventors' continuous study to establish a more economical and simple process for producing compounds of formula (1), a novel, straightforward and efficient process has been found enabling synthesis of compounds (1) in two steps from easily available 1,4-cyclohexanedione, by employing novel intermediates and a novel way of conversion of such new intermediates to compounds (1). Particularly, the process of our invention raises a potential to prepare the compound (1) enriched by a single enantiomer thereof.


According to the first aspect of the present invention, there is provided a process for preparing compounds of formula (1)



5

wherein R is hydrogen or acyl group, R₁ is hydrogen, lower alkyl or aralkyl group and R₂ is hydrogen,

comprising a reaction of a compound of formula (6),



10

wherein R is hydrogen or acyl group, R₃ and R₄ are either the same and each of them represents an alkoxy group of 1-4 carbons or they together form a C₂-C₅ alkylenedioxy group or an oxo- group ,

15 with an amine of general formula (7)




wherein R₁ is as hereinbefore defined,

20 in a presence of a reducing agent or a hydrogen gas with hydrogenation catalyst, optionally followed, if a compound (1) with R = acyl is produced, by hydrolysis of the acyl group to hydrogen, and isolation of the resulted compound of formula (1) as a free base or as an acid addition salt, incl. any hydrate or solvate thereof.

In a particular aspect, there is provided a process as defined above wherein the compound of formula (1) is produced substantially enriched by a single enantiomer, by using stereospecifically reducing agent or a chiral hydrogenation catalyst.

5 The compounds of formula (6) wherein R_3 and R_4 are as defined above are novel and they represent a second aspect of the present invention. As the preferred compounds, 2-amino-6-oxo-4,5,6,7-tetrahydrobenzothiazole (compound (6), $R = H$, R_3 , R_4 is =O group), 2-amino-6,6-dimethoxy-4,5,6,7-tetrahydrobenzothiazole (compound (6), $R = H$, R_3 and R_4 are methoxy groups), 2-acetamido-6-oxo-4,5,6,7-tetrahydrobenzothiazole (compound (6), $R = \text{acetyl}$, R_3 , R_4 is =O group) and 2-acetamido-6,6-dimethoxy-4,5,6,7-tetrahydrobenzothiazole (compound (6), $R = \text{acetyl}$, R_3 and R_4 are methoxy groups) are claimed, as well as acid addition salts thereof.


 In a third aspect, the present invention provides a process for producing a compound of formula (6) wherein R_3 and R_4 are as defined above, said process comprising
15 bromination of 1,4-cyclohexandione (5) in an alcoholic solvent to yield a compound of formula (8)



20 wherein R_3 and R_4 are as defined above,

followed by reaction with thiourea or N-acylthiourea. In a preferred aspect, the reaction is performed without isolation of intermediate product of the general formula (8). If desired, the acyl group R may be subsequently hydrolysed to hydrogen.

 The overall process of the present invention is illustrated by the
25 following reaction scheme:




As shown by the above reaction scheme, the first step of a process of the present invention is characterised by a novel finding that 1,4-cyclohexanedione (5) can be selectively monobrominated to give an intermediate (8). Though seeming to be a simple compound, no records of successful monobromination of 1,4-cyclohexanedione have been found in any prior art disclosure and the intermediate (8) has not been described and/or characterised so far.

We have found out that the monobromination of 1,4-cyclohexandione (5) cannot be achieved if attempting to brominate the substrate under "classical" conditions, e.g. by using bromine in acetic acid or in a halogenated hydrocarbon (e.g. in chloroform) and at slightly elevated temperatures, which conditions could be regarded by a skilled chemist as quite suitable for the reaction as the substrate is well soluble in these solvents. Indeed, bromine apparently reacts with the substrate, but the reaction product is different: immediate elimination of hydrogen bromide and formation of hydroquinone system follow the bromination.

After thorough experimental work resulting into our invention, we were successful in finding conditions, under which the desired compound (8) could be formed reliably. It was found that a substantial condition for successful monobromination of (5) is to perform the bromination reaction in an alcohol as a reaction medium. A relatively stable product of bromination can be obtained only under a condition if the keto-groups are masked as a ketal; the alcohol is thus also a reaction partner for such ketalization. As the alcohol, a lower alkanol of one to four carbon atoms (e.g. methanol, ethanol, n-propanol) or an alkylene glycol of two to five carbon atoms (e.g. ethylene glycol or 2,2-dimethylpropylene glycol) may be employed, though other alcohols are not specifically excluded. Methanol is the most preferred alcohol for the bromination.

The proper substrate for the bromination is thus in fact a compound of formula (9)




wherein R3 and R4 either represent the same alkoxy group or they together form an alkylendioxy group. The term "alkoxy" represents a straight or branched alkoxy group having one to four carbon atoms; the term "alkylendioxy" represents a group wherein the alkylene moiety consists from two to five carbon atoms.

The product of the bromination of the formula (8) is indeed still a quite unstable compound. Thus, the bromination reaction should advantageously be followed by the reaction with thiourea without isolation of the compound (8); however, the step of isolation of the intermediate (8) is not specifically excluded from this invention. The second advantage of the alcoholic solvent is that it can serve as a reaction medium also for the reaction with thiourea. The primary product (6) is the ketal (i.e. R₃,R₄ are alkoxy groups or alkylendioxy group) ; however, the ketal can be easily converted into a 6-oxo compound in the presence of water. Water can be either added intentionally, both after the reaction with thiourea (to the reaction mixture or to isolated ketal) and before the reaction of thiourea, or it may be present in the reaction mixture as a product of reaction from preceded steps. The deketalisation to 6-oxo compound generally requires the presence of an acid; herein, the hydrogen bromide liberated from the reaction with thiourea can serve as the acid catalyst.

It is not decisive whether the product (6) is isolated after the reaction with thiourea in a form of a ketal or as a 6-oxo compound or as a mixture of these two compounds. Both types of the substrate are equally suitable for the subsequent amination. The product (6) may be isolated as a free base or as an acid addition salt; preferred salt is a hydrobromide.

In the last (i.e. the second) production step, the isolated product (6) or the mixture of products (6) is subjected to the reductive amination with ammonia or corresponding alkylamine (7) in presence of a reducing agent to give requested compound of the formula (1). The reaction is advantageously carried out in a suitable solvent or in a mixture of solvents such as methanol, ethanol, tetrahydrofuran, dimethylformamide, conveniently at temperatures of between 0–50°C, preferably at temperatures of between 0–30°C. Sodium cyanoborohydride is the reducing agent of choice as it selectively reduces imines while it doesnot reduce oxo- group (so that the yet unreacted starting (6) is basically not reduced to an alcohol in a side reaction), although other conventional metal hydrides as sodium borohydride or diisobutylaluminium hydride can also be employed. Alternatively, the reaction may be performed under conditions of catalytic hydrogenation; however, care is to be taken in selection of hydrogenation catalysts as the sulphur present in the molecule of the substrate can poison the catalyst. Useful hydrogenation catalyst is e.g. palladium/carbon catalyst.


When using an alkylamine for reductive amination, an imine of the formula (10)




wherein R and R₁ are the same as above, is formed as an intermediate. Such compound may be isolated in case of need and can be subjected to a reduction to (1) in a separate step.

If using conventional reduction or hydrogenation agents, the desired compound (1) is formed as a racemate; however, in a specific feature of the invention, there may be employed also agents allowing enantioselective reductive amination, thus yielding a compound of formula (1) which is enriched by the desired enantiomer. Such agents are known in the art, e.g. as described in US 5,292,893 or WO 97/11934.

The reductive amination of a compound (6), particularly an enantioselective reductive amination, may be also performed indirectly. This alternative is useful for preparation of compounds of formula (1) wherein the substituent R_1 is an alkyl group of at least three carbon atoms. The reaction partner of the compound of the formula (6) is now a precursor of the amine (7) – a chiral amine of the formula (7a)



bearing a hydrogen-replaceable substituent X on the centre of chirality. Such an amine reacts with the compound (6) - under conditions substantially identical with those as described above- to yield a compound of the formula (11).



The presence of a centre of chirality in (7a) aids to direct the reductive amination asymmetrically so that the compound (11) may be preparable as enriched by one enantiomer.

In a subsequent step, the substituent X should be replaced by hydrogen to form the compound of formula (1), substantially without racemization on the tetrahydrobenzthiazole ring. Any substituent replaceable by hydrogen but resistant to the conditions of reductive amination may be chosen; advantageously, the starting chiral amine (7a) may be an amine wherein X is a hydroxy group or a halogen atom.

It is apparent that the substituent R_5 in the compounds (7a) and (11) is a precursor for the R_1 substituent and should be an alkyl group of two carbon less than

the desired group R_1 . Furthermore it is apparent that R_5 and X must not be identical to maintain the chiral centre.

The compounds of general formula (11), the process of their formation and that of their transformation to compounds (1) are novel and form another aspect of the present invention.

Another alternative of reductive amination is based on the reaction of a compound of formula (6) with N-alkylhydroxylamine. In this case, an iminoxide is formed as the intermediate and it may be reduced similarly as the imine. Reduction of the iminoxide may also produce a compound enriched by one enantiomer.

The enrichment, in optimum case, could be close to 100%. If not, the amount of undesired enantiomer in the product may be further diminished by means of conventional optical resolution, e.g. by fractional crystallisation of salts with an optically active acid or by chromatography on a chiral phase.

Before or after the reductive amination, the acyl group R may be hydrolysed to hydrogen, if desired. Such hydrolysis can be performed by conventional methods, preferably by alkaline hydrolysis.

If the product of the reductive amination (with or without a subsequent resolution into optical enantiomers) is a compound wherein R_1 is hydrogen, it may be converted into another compound of formula (1) wherein R_1 is an alkyl group or an aralkyl group, by conventional methods of alkylation or aralkylation of primary amines.

The compounds of formula (1) may be isolated from reaction mixture as free bases or they may be converted into acid addition salts and isolated in solid state by methods known per se. Preferred acids useful to form addition salts are pharmaceutically acceptable acids such as, e.g. hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, oxalic, maleic, fumaric, tartaric, malic, benzoic, methanesulfonic, benzenesulphonic or p-toluenesulfonic acid. The compounds or salts thereof may be also isolated in solvated or hydrated form.

Specifically, the process of our invention and the novel intermediates resulted therefrom are advantageous in the preparation of pramipexole. In the most preferred

arrangement for production of pramipexole, the bromination reaction of 1,4-cyclohexandione is carried out in methanol under temperatures close to ambient, and the reaction with thiourea with the product of bromination is performed without isolation thereof, i.e. in the same reaction vessel. The product of the condensation, either the ketal or
5 the ketone, is insoluble in the reaction medium and can be easily isolated by conventional methods of filtration or centrifugation, as a hydrobromide or as a free base. If necessary, this intermediate product may be further purified by conventional methods, e.g. by crystallisation.

In the following step, the above intermediate reacts with propylamine
10 and/or with a salt of propylamine, in a presence of a reducing agent such as sodium cyanoborohydride, in temperatures close to ambient and in an inert organic solvent. The intermediating imine is not isolated. After removal of inorganic salts and the excess of propylamine, the desired product can be isolated as a salt with an inorganic or organic acid, e.g. with hydrochloric acid, by crystallisation from a suitable solvent, e.g. from
15 ethanol.

Alternately, undirect reductive amination with a chiral amine as described above may be employed for the pramipexole synthesis.. The advantageous chiral amine for this reaction is S(-) 2-hydroxypropylamine. The intermediating product is 2'-hydroxypramipexole (compound (11), $R_5=CH_3$, $X=OH$) and, in the subsequent
20 step, the OH- group on the side chain thereof is removed by a suitable dehydroxylation agent or method, e.g via a iodo-compound (compound (11), $R_5=CH_3$, $X=J$).

If the produced pramipexole is a racemate or it is not sufficiently enriched by the desired S(-) enantiomer, there may be employed a step of optical resolution, i.e. the removal of the undesired R(+) enantiomer by a fractional
25 crystallisation of salts of both enantiomers of pramipexole with an optically active acid. Advantageously, L(+) tartaric acid is suitable for use in such resolution step.

The present invention is more particularly described and explained by the following examples. It is to be understood, however, that the present invention is not limited to these examples and various changes and modifications may be made without
30 departing from the scope of the present invention.

Examples

Example 1 Preparation of 2-amino-6-oxo-4,5,6,7-tetrahydrobenzothiazole [Compound (6), R= H, R₃+R₄ = oxo]

- 5 8.8 g of 1,4-cyclohexanedione was dissolved with 40 mL of methanol. Solution was cooled to 10°C and 4.2 mL of bromine was dropwise added at this temperature during 15 minutes. The temperature was kept at 10°C for one hour and then allowed to rise to 20°C. When bromine colour faded, 7.2 g of thiourea was added. The reaction mixture was then diluted with 20 mL of water and heated under reflux for 2 h.
- 10 White precipitate was removed by filtration and filtrate was concentrated in vacuo and cooled. The crystals of product (6) were filtered off and washed with a cold water.
- ¹H-NMR spectrum in D₂O: 2.12 ppm (t, 2 H) C(4); 2.59 ppm (t, 2H) C(5); 2.86 ppm (s, 2H) C(7) ; 4.76 ppm (solvent)

15 Example 2 Preparation of 2-amino-6,6-dimethoxy-4,5,6,7-tetrahydrobenzothiazole hydrobromide [Compound (6), R=H, R₃=methoxy, R₄ = methoxy]

- 30 g of 1,4-cyclohexanedione was dissolved in 140 mL of methanol. Solution was cooled to 10°C and 13.6 mL of bromine was dropwise added at this temperature during 15 minutes. The temperature was kept at 10 °C for one hour and
- 20 then allowed to rise to 20°C. When bromine colour faded, 24.5 g of thiourea was added. The reaction mixture was then stirred overnight at ambient temperature. After cooling down to 5°C, the precipitated crystals of the product were filtered off and washed with cold methanol. Yield: 54.8 g. For analytical purposes, a sample of the product was recrystallized from hot methanol.
- 25 ¹H – NMR spectrum in D₂O: 2.05 ppm (t, 2 H) C(4); 2.50 ppm (t, 2H) C(5) ; 2.79 ppm (s, 2H) C(7) ; 3.24 ppm (s, 6 H) CH₃O
- ¹³C – NMR in DMSO : C(4) 21.98 ppm , C(5) 28.02 ppm , C(7) 32.41 ppm , C(6) 100.8 ppm ; thiazole ring C(2') 164 ppm , C(4') 134 ppm, C(5') 112 ppm ; methoxy group 48.3 ppm.

Example 3 Preparation of 2-amino-6-oxo-4,5,6,7-tetrahydrobenzothiazole [
Compound (6), R= H, R₃+R₄= oxo]

21.9 g of the dimethoxy compound from example 2 was suspended in
5 205 ml of water and heated to 45°C for dissolution. Then 7.3 ml of concentrated HCl
was added and the solution was stirred at 60°C for 30 minutes. Reaction mixture was
cooled to ambient temperature and neutralised with 40% aqueous NaOH. Precipitated
solid was filtered off, washed with water and dried to yield 10.9 g of the title product.
For analytical purposes, the sample of the product was recrystallized from hot
10 isopropanol.

¹H – NMR in DMSO : 2.78 ppm (t , 2 H) C(4); 2.61 ppm (t , 2H) C(5) ; 3.37 ppm (s ,
2H) C(7)

¹³C – NMR in DMSO : C(4) 25.09 ppm , C(5) 38.87 ppm , C(7) 37.08 ppm , C(6)
207.07 ppm ; thiazole ring C(2') 167.1 ppm ; C(4') 144.08 ppm , C(5') 111.00 ppm

Example 4 Preparation of (±)2- amino-6-propylamino-4,5,6,7-tetrahydrobenzothia-
15 zole [Compound (1), R= hydrogen, R₁= propyl, R₂ = hydrogen]

3.0 g of propylamine hydrochloride was dissolved in 60 mL of methanol
and 2.2 g of 2-amino-6-oxo-4,5,6,7-tetrahydrobenzothiazole was added. 1.1 mL of
propylamine was poured into the mixture to set pH within the range 6.5–7.0. Then 0.7 g
of sodium cyanoborohydride was added and the reaction mixture was stirred for 70
20 hours under nitrogen atmosphere and at ambient temperature. The reaction mixture was
diluted with 50 mL of water and 10 mL of conc. hydrochloric acid. Solvents were
removed by evaporation, the yellow residue was stirred with ethanol and the solid was
removed by filtration. The filtrate was made alkaline with potassium hydroxide solution
and the precipitate was again removed by filtration. Filtrate was several times
25 evaporated with ethanol to remove propylamine. The residue was dissolved in hot
ethanol and resulting crystals were collected and dried. Reaction product was
characterised by NMR spectra.

¹H – NMR spectrum in CDCl₃ : 0.93 ppm (t , 3 H) C(3'); 1.51 ppm (sextet , 2 H) C(2'); 2.65 (t , 2H) ppm (s) C(1'); 2.6 ppm (m, 2H) C(4); 1.99 ppm (m , 1H) + 1.70 ppm (m , 1H) C(5); 2.98 ppm (m , 1H) C(6); 2.82 ppm (dd , 1H) + 2.40 ppm (m , 1H) C(7);
¹³C – NMR in CDCl₃: C(4) 29.47 ppm , C(5) 24.92 ppm , C(7) 30.1 ppm , C(6) 53.92 ppm , C(1') 49.20 ppm , C(2') 23.48 ppm , C(3') 11.76 ppm ; thiazole ring C(2') 165.49 ppm ; C(4') 145.02 ppm , C(5') 116.31 ppm

Example 5 Preparation of 2-acetamido-6-oxo-4,5,6,7-tetrahydrobenzothiazole hydrobromide [Compound (6), R= acetyl, R₃+R₄ = oxo]

8.8 g of 1,4-cyclohexanedione was dissolved in 40 ml of methanol. The solution was cooled on an ice bath to 10°C and 4.4. ml of bromine was added dropwise in 10 minutes. The solution was stirred at the same temperature for 1.5 hours and then the temperature was allowed to rise to ambient. After the colour disappeared, 11.1 g of N-acetylthiourea and 20 ml of water were added and the solution was heated under reflux for 2 hours. Hot reaction mixture was filtered, the filtration cake was washed with hot methanol and the combined filtrates were heated to reflux with activated charcoal. The hot suspension was filtered and the filtrate was concentrated in vacuo to yield 3.4 g of a solid product.

Example 6 Preparation of pramipexole dihydrochloride from (±)2-amino-6-propylamino-4,5,6,7- tetrahydrobenzothiazole

5 g of (±)2-amino-6-propylamino-4,5,6,7- tetrahydrobenzothiazole was dissolved in 25 ml of methanol and 4.0 g of L(+)-tartaric acid was added under stirring. The mixture was cooled and the white crystals were filtered off, washed with methanol and dried to yield 4.8 g of pramipexole tartrate. The crystals were dissolved in 10 ml of water and concentrated hydrochloric acid was added to dissolve the solid. The solution was cooled to 10 °C and a solution of 10 g of potassium hydroxide in 10 ml of water was added. The mixture was stirred for 30 minutes and the white crystals were removed by filtration and washed with cold water. The crystals were redissolved in 7 ml of ethanol and gaseous hydrogen chloride was bubbled through the solution at 10 °C. After 1 hour of stirring, the

precipitated solid was separated by filtration, washed with cold ethanol and dried to yield 2.1 g of white crystals. The product is identical with pramipexole dihydrochloride by NMR and has optical rotation $[\alpha]_{\text{D}}^{20} = -48.78^\circ$ ($c=1$, MeOH)

5 Example 7 Preparation of 2,6 - diamino - 4,5,6,7 - tetrahydro - 1,3 - benzothiazole
[Compound (1), R=R₁=R₂ = hydrogen]

A solution of 2 - amino - 6 -oxo - 4,5,6,7 - tetrahydro - 1,3 - benzothiazole (2.0 g , 12 mmol) , ammonium acetate (9.2 g , 120 mmol) and sodium cyanoborohydride (0.53 g , 8.4 mmol) in 40 mL of methanol was stirred under nitrogen at ambient temperature for 48 hours . Concentrated HCl was added and methanol was removed in vacuo. The residue was mixed with water and solution was made alkaline with NaOH. White precipitate was removed by filtration , washed with water and dried to give 1.0 g of the title compound. The structure of the product was confirmed by NMR.

Example 8 Preparation of 2-amino-6-propylamino-4,5,6,7-tetrahydrobenzothiazole
using S(+) 1-amino-2-propanol

Step 1) Preparation of (2S)-1-(2'-amino-4',5',6',7'-tetrahydro-benzothiazol-6'-ylamino)-propan-2-ol [Compound (1), R= H, R1=2(S)-hydroxypropyl , R2 = hydrogen]

6.7 g of S-(+)-1-amino 2-propanol was dissolved in 125 mL of methanol and 10 g of 2-amino-6-oxo-5,6,7,8-tetrahydrobenzthiazole was added under nitrogen. The yellow suspension was stirred at 5-7 C for 20 minutes and 3,85 g of sodium cyanoborohydride was added. After 20 hours of stirring the reaction mixture was acidified with hydrochloric acid and the solvent was removed in vacuo. Crystalline residue was dissolved in water and water solution was alkalisied with 25 % aqueous NaOH. Precipitated solid was filtered off, washed with cold water and dried to yield 5.5 g of the title product (m.p. 75 - 85°C).

^1H – NMR in DMSO : 1.14 ppm (d , 3H) C(3'); 4.12 ppm (m , 1H) C(2') ; C(1') 3.0 ppm (m, 2H) ; 3.54 ppm (m , 1H) C(6) ; 3.10 ppm (m , 2H) C(7); 2.34 ppm (m , 2H) C(3) ; 2.09 ppm (m , 2H) C(4)

^{13}C – NMR in DMSO : C(4) 23.2 ppm , C(5) 20.9 ppm , C(7) 24.69 ppm , C(6) 52.65 ppm , C(1') 51.51 , C(2') 62.30 , C(3') 21.02 ppm ; thiazole ring C(2') 168.7 ppm ; C(4') 132.8 ppm , C(5') 110.83 ppm

Step 2 : Preparation of (2'S) - N-6-(2'-Iodo-propyl)-4,5,6,7-tetrahydro-benzothiazole-2,6-diamine dihydroiodide [Compound (1), R= H, R1=2(S)-iodopropyl , R2 = hydrogen]

5.0 g of the product of Step 1, 5.0 g of red phosphorus and 150 mL of 57 % hydroiodic acid was refluxed for 25 hours. Reaction mixture was filtered and the filtrate was evaporated in vacuo. The crystalline residue was stirred with acetone , crystals removed by suction and washed with acetone to give 6.5 g of the title product with m.p. 213-222C.

^1H – NMR in DMSO : 2.0 ppm (d , 3H) C(3'); 4.45 ppm (m , 1H) C(2') ; 3.62 ppm (m, 2H) C(1') ; 2.80 ppm (m , 1H) C(4) ; 2.1 - 2.5 ppm (m , 2H) C(5) ; 3.87 ppm (m , 2H) C(6); 3.2 ppm (m , 2H) C(7)

^{13}C – NMR in DMSO : C(4) 17.42 ppm , C(5) 17.17 ppm , C(7) 26.09 ppm , C(6) 56. ppm , C(1') 55.54 , C(2') 26.90 ppm , C(3') 22.42 ppm ; thiazole ring C(2') 171.0 ppm, C(4') 134.34 ppm , C(5') 112.88 ppm

Step 3 : Preparation of 2-amino-6-propylamino-4,5,6,7- tetrahydrobenzothiazole

6.0 g of the product of Step 2 was dissolved in 100 mL of water acidified with 1 mL of 36 % hydrochloric acid. 0.3 g of 10 % Pd/C catalyst was added and the reaction mixture was hydrogenated at 100 $^{\circ}\text{C}$ and 4.0 MPa of hydrogen for 8 hours. Catalyst was filtered off , washed with water and the filtrate was made alkaline with addition of 50 % sodium

hydroxide solution. Precipitated product was filtered off, washed with water and air-dried to give 1.8 g of the title product with m.p. 145 - 150 °C and optical purity 42 %..

Example 9 Preparation of (±) 2-amino-6-propylamino-4,5,6,7-tetrahydro-benzothiazole [Compound (1), R= hydrogen, R₁= propyl, R₂ = hydrogen] dihydrochloride

A 150 mL Paar autoclave was charged with 3.5 g of 2-amino-6-oxo-4,5,6,7-tetrahydrobenzthiazole, 50 mL of tetrahydrofuran, 2.0 mL of propylamine and 0.8 g of 10%Pd/C catalyst. The reaction mixture was heated at 100°C, 40 bars of hydrogen and 400 RPM for 48 h. Catalyst was removed by filtration, filtrate evaporated to dryness, the residue dissolved in ethanol and solution saturated with gaseous hydrogen chloride. Precipitated crystals were removed by suction, washed with ethanol and dried to give crystalline product with m.p. 268 - 278 °C.

Example 10 Preparation of (+) 2 - amino - 6 - oxo - 4,5,6,7 - tetrahydrobenzothiazole
[Compound (6), R= H, R₃+R₄ = oxo] hydrobromide

340 g of 2-amino-6,6-dimethoxy-4,5,6,7-tetrahydrobenzothiazole hydrobromide (Example 2) was mixed with 290 mL of hot water. The mixture was evaporated under reduced pressure at low temperature (50-55°C), and the residue was three times evaporated with 170mL of acetone. The residue was mixed with 340 mL of acetone and after 1 hour of stirring the solid was filtered off and air dried to give 168 g of the title product.

Example 11 Preparation of (+)2 - amino - 6 - propylamino - 4,5,6,7 - tetrahydro
benzothiazole


10 g of 2 - amino - 6 - oxo - 4,5,6,7 -tetrahydrobenzthiazole hydrobromide was
5 added to the mixture of 5 mL of propylamine and 125 mL of methanol under
nitrogen. The mixture was stirred for 20 minutes. Then 2,7 g of sodium
cyanoborohydride was added in parts during 30 minutes . The solution was acidified
with 2mL of acetic acid to the value pH=7-8. Reaction mixture was worked up after 20
hours of stirring by addition of 8 mL of HCl. Precipitated solid was removed by suction
10 and the filtrate was evaporated to dryness. The solid residue was heated at reflux with
250mL of ethanol for 20 minutes. Undissolved part was filtered off and the hot filtrate
was concentrated under reduced pressure and cooled on an ice bath. Precipitated
crystals were removed by suction and the cake was washed with cold ethanol to give 5.1
g of a product with m.p. 245-257°C .

15 The solid product was dissolved in 28 mL of of water and the solution was
alkalinized with 25% aqueous NaOH. The mixture was stirred for 2 hours and
precipitated crystals were removed by suction and washed with water to give 2.7 g of
the title product with m.p. 147-149°C.

CLAIMS

1. A compound of the general formula (6)

5



wherein R is hydrogen or acyl group, R₃ and R₄ are either the same and each of them represents an alkoxy group of 1-4 carbons or they together
10 form an C₂-C₅ alkylenedioxy group or an oxo-group ,
and acid addition salts thereof.

2. A compound according to claim 1 wherein R is hydrogen or acetyl group and R₃ and R₄ each represent methoxy, ethoxy or n-propoxy group.

3. A compound according to claim 1 wherein R₃ and R₄ together form
15 an oxo group.

4. A compound according to any one of precedent claims selected from the group consisting of 2-amino-6-oxo-4,5,6,7-tetrahydrobenzthiazole, 2-amino-6,6-dimethoxy-4,5,6,7-tetrahydrobenzothiazole, 2-acetamido-6-oxo-4,5,6,7-tetrahydrobenzothiazole and 2-acetamido-6,6-dimethoxy-4,5,6,7-tetrahydrobenzthiazole, and acid addition salts thereof.


20 5. A process for preparation of a compound of formula (6) as defined in any one of claims 1 to 4, which comprises bromination of 1,4-cyclohexanedione by bromine in an alcoholic solvent, followed by treatment of the reaction mixture with thiourea or N-acylthiourea and isolation of the compound of formula (6) from the reaction mixture.

6. A process according to the claim 5 wherein the alcoholic solvent is selected from the group consisting of methanol, ethanol, n-propanol, ethylene glycol or 2,2,-dimethylpropyleneglycol.

7. A process according to the claim 6 wherein the solvent is methanol.

5 8. A process according to any of claims 5-7 wherein the reaction mixture is treated with thiourea.


9. A process of preparation of a compound of formula (1)



10

wherein R is hydrogen or acyl group, R1 is hydrogen, lower alkyl or aralkyl group and R2 is hydrogen,

comprising reacting a compound of formula (6)



15

wherein R is hydrogen or acyl group, R3 and R4 are either the same and each of them represents an alkoxy group of 1-4 carbons or they together form an C2-C5 alkylendioxy group or an oxo- group ,


20 A) with an amine of the formula (7)



wherein R_1 is as hereinbefore defined,

under presence of a reduction agent or a hydrogen gas with hydrogenation catalyst or,
alternately, isolating the imine intermediate and reducing it by reaction with said
5 reduction agent or by hydrogenation or,


B) reacting said compound of formula (6) with a chiral amine of formula (7a)



10 wherein R_5 is an alkyl group of two carbon less than R_1 , and X is a hydrogen-replaceable leaving group not identical with R_5 ,

under presence of a reduction agent or a hydrogen gas with hydrogenation catalyst or,
alternately, isolating the imine intermediate and reducing it by reaction with said
reduction agent or by hydrogenation, to yield a compound of formula (11)

15



followed by a step of substitution of the group X in the compound of formula (11) for hydrogen.

20 10. A process according to claim 9 wherein the reduction agent is a metal hydride.

11. A process according to claims 9 and 10 wherein the reduction agent is an agent allowing forming a product enriched with one enantiomer thereof.

12. A process according to claims 9-11 further comprising a step of hydrolysis of the acyl group to hydrogen, when a compound (1) with R = acyl is produced by the said process.

5 13. A process according to claims 9-12 wherein in the chiral amine of formula (7a) R is hydrogen, R₅ is methyl and X is hydroxy or halogen group.

14. A process according to claims 9-13 wherein the product obtained is a mixture of optical isomers enriched with one of the optical isomers.

15 15. A process according to claims 9-14 further comprising a step of resolution of the obtained racemic product into single optical enantiomers thereof.

16. A process according to claim 9-15 further comprising a step of isolation of the resulted compound of formula (1) from the reaction mixture as a free base or as an acid addition salt.

15 17. A process according to claims 9-16 wherein the isolated product is substantially S(-)-2-amino-6-propylamino-4,5,6,7-tetrahydrobenzo-thiazole (pramipexole) or its acid addition salts.

18. A process according to claims 16-17 wherein the product is isolated as the dihydrochloride.

20 19. A process according to any one of claims 9-18 wherein the starting compound of formula (6) is obtained following the process of claims 5-8.


20. A process for preparation of 2-amino-6-propylamino-4,5,6,7-tetrahydrobenzothiazole according to claims 9-11 and 14-17, comprising the step of reaction of propylamine and/or its salt with a compound selected from a group consisting of 2-amino-6-oxo-4,5,6,7-tetrahydrobenzothiazole, 2-amino-6,6-dialkoxy-4,5,6,7-tetrahydrobenzo-thiazole or a mixture of both compounds, under presence of a
25 reduction agent or a hydrogen gas with hydrogenation catalyst or, alternatively,

isolating the imine intermediate and reducing it by reaction with said reduction agent or by hydrogenation.

21. A process according to claim 20 further comprising the step of resolution 2-amino-6-propylamino-4,5,6,7-tetrahydrobenzothiazole into its enantiomers using L(+) tartaric acid as the resolution agent and isolation the enantiomer.

22. The process according to claim 21 wherein the isolated product is substantially S (-)-2-amino-6-propylamino-4,5,6,7-tetrahydrobenzothiazole (pramipexole) or its acid addition salt.

23. A compound of formula (11)



wherein R is hydrogen or acyl group, R₅ is a lower alkyl group and X is a hydrogen-replaceable leaving group not identical with R₅.

24. A compound according to claim 23 wherein R is hydrogen, R₅ is methyl and X is hydroxy or halogen group.

25. Use of 1,4-cyclohexandione for production of compounds of formula (6) as defined in claims 1 - 4.

26. Use of a compound of formula (6) as defined in claims 1 - 4 for production of compounds of formula (1) as defined in claim 9.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/NL 01/00680

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D277/82 //C07M7:00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,Y	EP 0 251 077 A (BOEHRINGER INGELHEIM KG) 7 January 1988 (1988-01-07) the whole document, particularly example 1	1-24
X,Y	KOZIKOWSKI A P ET AL: "Synthesis of the thiazolone analogue of the acetylcholinesterase inhibitor, huperzine A" HELVETICA CHIMICA ACTA, vol. 77, no. 5, 10 August 1994 (1994-08-10), pages 1256-1266, XP002186318 the whole document, particularly page 1258, scheme 1 --- -/--	1-24

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

21 December 2001

Date of mailing of the international search report

18/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Aillard, M

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/NL 01/00680

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 186 087 A (DR. KARL THOMAE GMBH) 2 July 1986 (1986-07-02) cited in the application the whole document	1-24
Y	----- SCHNEIDER C S ET AL: "Dopamine autoreceptor agonists: resolution and pharmacological activity of 2,6-diaminotetrahydrobenzothiazole and aminothiazole analogue of apomorphine" JOURNAL OF MEDICINAL CHEMISTRY, vol. 30, no. 3, March 1987 (1987-03), pages 494-498, XP002186199 cited in the application the whole document -----	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 01/00680

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 251077	A	07-01-1988	DE 3620813 A1	23-12-1987
			AT 70060 T	15-12-1991
			AU 593357 B2	08-02-1990
			AU 7457887 A	24-12-1987
			CA 1322372 A1	21-09-1993
			CS 8704511 A3	13-05-1992
			DD 259623 A5	31-08-1988
			DE 3774942 D1	16-01-1992
			DK 168437 B1	28-03-1994
			EP 0251077 A1	07-01-1988
			ES 2040226 T3	16-10-1993
			FI 872719 A ,B,	22-12-1987
			GR 3003656 T3	16-03-1993
			HU 47091 A2	30-01-1989
			IE 60676 B	10-08-1994
			JP 63017872 A	25-01-1988
			KR 9504000 B1	22-04-1995
			NO 872574 A ,B,	22-12-1987
			NZ 220800 A	21-12-1989
			PH 24207 A	10-04-1990
			PT 85100 A ,B	01-07-1988
			SU 1494867 A3	15-07-1989
			US 4981862 A	01-01-1991
			ZA 8704470 A	22-02-1989
EP 186087	A	02-07-1986	DE 3447075 A1	03-07-1986
			DE 3508947 A1	18-09-1986
			AT 45735 T	15-09-1989
			AU 583874 B2	11-05-1989
			AU 5154485 A	17-07-1986
			BG 62023 B2	30-12-1998
			BR 1100678 A3	13-10-1999
			CA 1263653 A1	05-12-1989
			CS 9104099 A3	16-09-1992
			DD 242230 A5	21-01-1987
			DE 3572485 D1	28-09-1989
			DK 590285 A	23-06-1986
			EP 0186087 A1	02-07-1986
			ES 550235 D0	16-01-1987
			ES 8702787 A1	01-04-1987
			ES 556873 D0	01-08-1987
			ES 8707513 A1	16-10-1987
			ES 556874 D0	01-08-1987
			ES 8707514 A1	16-10-1987
			ES 556875 D0	01-08-1987
			ES 8707515 A1	16-10-1987
			FI 855102 A ,B,	23-06-1986
			GR 853126 A1	22-04-1986
			HK 78692 A	23-10-1992
			HU 39736 A2	29-10-1986
			IE 58863 B	17-11-1993
			IL 77415 A	19-03-1990
			JP 1854941 C	07-07-1994
			JP 61155377 A	15-07-1986
			JP 5072907 B	13-10-1993
			KR 9309791 B1	11-10-1993
			LU 90208 A9	06-04-1998
			MX 9202792 A1	30-06-1992

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/NL 01/00680

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 186087	A	NO 855195 A ,B,	23-06-1986
		NZ 214661 A	26-04-1990
		PH 24533 A	03-08-1990
		PT 81735 A ,B	01-01-1986
		SG 82492 G	04-12-1992
		US 4843086 A	27-06-1989
		US 4886812 A	12-12-1989
		US 4731374 A	15-03-1988
		ZA 8509731 A	26-08-1987
<hr/>			